



Available online at
ScienceDirect
www.sciencedirect.com

Elsevier Masson France
EM|consulte
www.em-consulte.com



Practice guidelines

French Guidelines For the Emergency Management of Headaches



X. Moisset^{a,b,*}, J. Mawet^c, E. Guegan-Massardier^d, E. Bozzolo^e, V. Gilard^f,
 E. Tollard^g, T. Feraud^h, B. Noëlleⁱ, C. Rondet^j, A. Donnet^{a,k}

^a Inserm U-1107, NeuroDol, Clermont Université, Université d'Auvergne, 49, boulevard François-Mitterrand, 63000 Clermont-Ferrand, France

^b CHU Gabriel Montpied, Service de Neurologie, Clermont Université, Université d'Auvergne, Clermont-Ferrand, France

^c Centre d'urgences céphalées, département de Neurologie, GH Saint-Louis-Lariboisière, Assistance Publique des Hôpitaux de Paris AP-HP, Université Paris Denis Diderot et DHU NeuroVasc Sorbonne Paris-Cité, Paris, France

^d Service de neurologie, hôpital Charles-Nicolle, 1, rue de Germont, 76000 Rouen, France

^e Service de neurologie, Pôle des Neurosciences Cliniques, CHU de Nice, Nice, France

^f Service de neurochirurgie, hôpital Charles-Nicolle, 1, rue de Germont, 76000 Rouen, France

^g Service de neuroradiologie, hôpital Charles-Nicolle, 1, rue de Germont, 76000 Rouen, France

^h Service d'accueil des urgences, hôpital Timone, boulevard Jean-Moulin, 264, rue Saint-Pierre, 13385 Marseille, France

ⁱ Cabinet privé, 35, allée de Champrond, 38330 Saint-Ismier, France

^j Faculté de médecine, Service de médecine générale, Université Pierre-et-Marie-Curie Paris 06, Paris, France

^k Centre d'évaluation et de traitement de la douleur, hôpital Timone, boulevard Jean-Moulin, 264, rue Saint-Pierre, 13385 Marseille, France

INFO ARTICLE

Historique de l'article :

Reçu le 6 mars 2016

Accepté le 8 juin 2016

Disponible sur Internet le

1 July 2016

1. Preamble

1.1. Requesting body

These guidelines were elaborated at the request of the French society for the study of migraine and headache disorders (SFEMC¹) and the French Society of Neurology (SFN²).

1.2. Topic

These guidelines concern recommended management practices for patients with headache disorders seen in an emergency setting. Four clinical aspects are distinguished: key elements of history taking and the physical examination, diagnostic and therapeutic strategies.

* Corresponding author. Inserm U-1107, NeuroDol, Clermont Université, Université d'Auvergne, 49, boulevard François-Mitterrand, 63000 Clermont-Ferrand, France.

E-mail address: xavier.moisset@gmail.com (X. Moisset).

¹ SFEMC Société Française d'Etude des Migraines et des Céphalées.

² SFN Société Française de Neurologie.

<http://dx.doi.org/10.1016/j.neurol.2016.06.005>

0035-3787/© 2016 Elsevier Masson SAS. All rights reserved.

These guidelines do not concern the management of patients with headache disorders seen outside the emergency setting. Specific recommendations are available for migraine, chronic daily headache, or cluster headache.

1.3. Patients concerned

These guidelines concern adult patients.

1.4. Professionals concerned

These guidelines are written for healthcare professionals involved in the management of patients with headache disorders seen in the emergency setting.

1.5. Guideline grade and methodology

These guidelines propose grade A, B or C recommendations according to the following modalities:

- Grade A recommendations are based on scientific evidence with a high level of proof, e.g. high-power randomized comparative trials without major bias and/or meta-analyses of randomized comparative trials, decision analysis based on well-conducted studies;
- Grade B recommendations are based on scientific conjecture issuing from intermediary level-of-proof studies, e.g. intermediate-power randomized comparative trials, well-conducted non-randomized comparative trials, cohort studies;
- Grade C recommendations are based on studies with a low level-of-proof studies, e.g. case-control studies, series of cases

Unless specified, the proposed guidelines are based on professional agreement among the members of the working group.

The lack of a level of proof does not imply that the proposed guidelines are not pertinent and useful. The lack of proof should incite the development of complementary studies when possible.

These guidelines were elaborated by the SFEMC and the SFN, in application of the methodology for the elaboration of good clinical practices (GCP). The pilot group was composed of six members: Anne Donnet (neurologist), Anne Revol (neurologist), Pierrick Giraud (Neurologist), Dominique Valade (emergency headache specialist), Pierre Michelet (intensivist), Philippe Cornet (general practitioner). The working group was composed of nine members: Xavier Moisset (neurologist), Jérôme Mawet (neurologist), Evelyne Guegan-Massardier (neurologist), Eric Bozzolo (neurologist), Vianney Gilard (neurosurgeon), Eléonore Tollard (neuroradiologist), Thierry Feraud (intensivist), Bénédicte Noëlle (neurologist) and Claire Rondet (general practitioner). It was headed by Anne Donnet. Xavier Moisset wrote the guidelines.

A reading group composed of members of the SFEMC and the SFN as well as independent healthcare professionals from these societies (general practitioners, intensivists, neuroradiologists) was consulted.

2. Introduction

Headache during the last three months is observed in approximately 15% of the general population [1]. Consequently, about 1% of patients seeking ambulatory care do so for headache [2], and 2% of patients attending an emergency room do so for headache [3,4]. Despite the fact that 95% of these patients return home with a diagnosis of benign primary headache, the attending physician must be able to detect secondary and potentially dangerous headache that requires referral [3,4].

Headache disorders are defined according to the criteria proposed in the 3rd version of the International Classification of Headache Disorders [5].

The present guidelines concern all types of headache disorders, whether primary or secondary, occurring as an isolated symptom leading an adult patient to seek emergency medical care (call to primary care physician, emergency squad or emergency room). There are different reasons for which patients seek emergency care for headache disorders:

- sudden-onset headache;
- first-occurrence headache or unusual headache not corresponding to prior experience;
- headache unresponsive to usual treatment;
- chronic headache no longer tolerated.

As a preamble, we recall that headache intensity and severity are not correlated. In the majority of cases—even for patients seen in an emergency setting—headache is a primary disorder. Primary headache can be highly disabling but does not in itself constitute a functional or vital risk. What is important is to identify secondary headache in order to institute adapted—sometimes emergency—treatment.

Until evidence to the contrary, sudden-onset and/or unusual headaches should be considered as secondary disorders warranting emergency complementary exploration.

Headache that the patient recognizes as usual, but with greater intensity and unresponsive to usual pain relievers, is generally a primary headache refractory to habitual treatment. In this situation, complementary exploration is not useful and—under the condition that the physical examination, notably the neurological examination, is normal—management should focus on achieving pain relief.

3. How can the four clinical presentations—recent onset-headache, recent progressive headache, recurrent paroxysmal headache, and chronic daily headache—be distinguished?

In daily practice, patients seek emergency care for four main types of headache disorders (Fig. 1):




Fig. 1 – Schematic representation of the four clinical presentations.

- recent sudden-onset headache;
- recent progressive headache;
- recurrent paroxysmal headache;
- chronic daily headache.

Different etiologies are grouped together in each of these four clinical presentations. These categories are nevertheless useful for guiding the search for the main causes of the patient's symptoms. The diagnostic approach will be rapidly defined by history taking, from the patient or accompanying persons. It is important to differentiate the characteristic features of the headache for which the patient seeks emergency care from headaches the patient has experienced in the past.

4. What are the key elements to search for in the patient's history?

In addition to the usual questions designed to recall the patient's past medical history and treatments, a small number of questions will provide a rapid first-line evaluation of the potential gravity of the patient's headache. This minimal check-list should be used for emergency room triage (or for telephone or face-to-face interrogation by the primary care physician or intensivist) (Professional agreement).

- When did the pain start? (recent disorder)
- How long did it take for the pain to reach peak intensity? (sudden-onset disorder)
- Have you already had this kind of pain? (unusual disorder)
- What were you doing when the pain started? (exercise...)
- Are there other symptoms: fever, photo- or phonophobia, vomiting, neck pain, focal neurological deficit, slow psycho-motor response, girdle pain?

- Is there a particular context: new medication, toxic substances, exposure to carbon monoxide; trauma; pregnancy or post-partum, cancer, systemic disease (including HIV); maneuver with dural injury or risk of injury within the preceding month (lumbar puncture, peridural analgesia, spinal nerve injections)...?
- Does body position affect the pain (relief or worsening in the supine position)?

5. How should the physical examination be conducted?

If an attending physician is present, a systematic search should be conducted to identify (Professional agreement):

- impaired vigilance;
- fever;
- hypertension;
- meningeal syndrome;
- focal neurological deficit (motor or sensorial deficit, diplopia, pupil anomaly, cerebellar syndrome);
- disorders affecting the eyes, sinuses, ears or the oral cavity that could explain the headache.

6. What first-line measures are required?

If secondary headache warranting emergency treatment is suspected, the patient should be referred to the nearest emergency ward in order to undergo complementary exploration (brain imaging, cerebrospinal fluid [CSF] analysis, blood tests). Ideally, the patient should be referred to a hospital with a neurology unit (Professional agreement).

Thus, EMERGENCY CARE is recommended for all patients presenting:

- sudden-onset or thunderclap headache (peak intensity within one minute);
- recent-onset or recently worsening (< 7 days) unusual headache;
- headache associated with fever (without other obvious general cause of fever such as a flu-like syndrome in an outbreak context);
- headache associated with neurological signs;
- headache suggestive of intoxication (particularly CO);
- headache in a context of immune deficiency.

In the case of telephone contact, the patient should be advised to seek medical assistance rapidly (primary care or on-call physician) in the following situations:

- the patient is familiar with this type of headache but it is unresponsive to usual pain relievers;
- the headache is semi-recent (8 days to 6 months) with no sign of worsening in the last few days;
- longstanding headache (> 6 months) with an intensity or frequency the patient cannot cope with.

History taking should also identify the underlying pattern of the headache: “Do you have headaches daily, permanently, or by acute attacks?”

If the patient reports acute attacks, a precise description (duration, localization, intensity, associated signs, triggering factors, number of headache days per month, treatments tried and their efficacy) will help determine the type of headache. A given individual may present several types of primary headache (for instance, migraine without aura and tension-type headache).

If the patient has chronic daily headache, the reported description will guide the diagnosis. Nevertheless, any recent change in the presentation of a primary headache is suggestive of a secondary headache, to be ruled out or not. Moreover, it is important to search for overuse of pain relievers (medication overuse). In all cases, chronic daily headache existing for less than six months should be explored (Professional agreement) [6].

In patients reporting primary headache, complementary explorations are not useful if the attacks are well known and longstanding, fulfilling all the diagnostic criteria [5].

Outside acute episodes, the general and neurological examination is normal in patients with primary headache (excepting the Claude-Bernard Horner sign that is sometimes observed between attacks in a context of cluster headache).

7. What are the specific clinical features of the main primary headache disorders?

7.1. Migraine, tension-type headache and cluster headache

The clinical criteria are described in the ICHD-3 [5]. The main elements are summarized in Table 1. For patients presenting a first cluster headache, brain magnetic resonance imaging (MRI), and MRI or computed tomography (CT) angiography of the head and neck should be performed (Professional agreement) [7].

7.2. Status migrainosus

In patients with a known migraine, status migrainosus corresponds to a persistent disabling attack lasting more than 72 h. Periods of relief lasting less than 12 h (following medication or during sleep) can occur. This is a rare entity [8]. Secondary headache must be ruled out before establishing the diagnosis.

7.3. Chronic daily headache

Chronic daily headache occurs in patients who previously had an underlying primary headache [6]. These headaches can be secondary to medication overuse, defined as use at least 15 days per month for at least three months of non-opioid pain relievers (acetaminophen, aspirin, non-steroidal anti-inflammatory drugs [NSAID]), or use 10 days per month of opioid, triptan-ergotamine pain relievers, and/or a combination of several medications [5].

8. Diagnostic and therapeutic management

8.1. Diagnostic management

8.1.1. Sudden-onset headache

8.1.1.1. Clinical signs.

Until evidence to the contrary, all patients complaining of headache who fulfill one criterion of the Ottawa clinical decision rule (Box 1) should be suspected of having subarachnoid hemorrhage (Perry et al., 2013) (Grade B).

Headache is the only clinical sign in one-third of patients with subarachnoid hemorrhage [9]. There are no supplementary clinical criteria providing evidence that complementary exploration can be postponed (Grade C) [10]. The Ottawa clinical decision rule can be applied for all cases of non-trauma headache not associated with neurological deficit, a past history of aneurysm, subarachnoid hemorrhage, brain tumor, or recurrent headache (≥ 3 episodes in ≥ 6 months) [11,12].

Table 1 – Main clinical features for the diagnosis of migraine, tension-type headache and cluster headache.

	Migraine	Tension-type headache	Cluster headache
Duration	4–72 h	30 min–7 days	15 min–3 h
Site	Generally unilateral	Bilateral	Orbito-temporal unilateral
Intensity	Moderate to severe	Mild to moderate	Very severe
Type	Often pulsating	Compression, band-like pressure	Boring, squeezing
Accompanying signs	Nausea, vomiting, photo- and phono-phobia	No nausea or vomiting, Photo- or phono-phobia possible	Homolateral autonomic signs, agitation
Impact of exercise	Aggravation	No change	No change
Number of prior attacks	≥ 5	≥ 10	≥ 5

Box 1.

The Ottawa clinical decision rule

Exploration to search for subarachnoid hemorrhage is warranted in all patients presenting at least one of the six criteria. This rule is for patients aged over 15 years who present with severe non-traumatic headache that reached peak intensity in less than one hour. It does not apply to cases with neurological deficit, history of aneurysm, subarachnoid hemorrhage, brain tumor, or recurrent headache (≥ 3 episodes in ≥ 6 months).

- 1- Age ≥ 40 years
- 2- Neck pain or stiffness
- 3- Witnessed loss of consciousness
- 4- Onset during exertion
- 5- Thunderclap headache (intensity $> 7/10$ in less than 1 min)
- 6- Limited neck flexion

The rule has an excellent negative predictive value, but poor specificity [11,12]. However, complementary emergency exploration may be warranted in certain cases of headache not meeting any of the Ottawa rule criteria.

8.1.1.2. Complementary explorations.

A patient who presents with sudden-onset headache or headache associated with a neurological deficit should have an emergency CT scan (Grade B) (Edlow et al., 2008). Cerebral CT angiography should also be performed in all patients in order to explore the arterial and the venous networks (Professional agreement). MRI with MR angiography can be the first-line exploration if therapeutic management is not delayed (Professional agreement).

Positive diagnosis of subarachnoid hemorrhage is confirmed by the cerebral CT scan without contrast injection that has 98% sensitivity when performed within 12 h of symptom onset (sensitivity falls to 93% after 24 hr, 85% at 5 days and 50% at 7 days) (Grade B) [5,13,14]. Although a CT scan without contrast injection is indispensable, it is not sufficient. The Guide for good use of medical imaging explorations published by the French Society of Radiology and the French Society of Nuclear Medicine under the auspices of the French

health authorities (HAS³) and the Nuclear safety authorities (gbu.radiologie.fr) recommends that a cerebral CT angiography should also be performed in all cases if no bleeding is visualized spontaneously, in order to explore the arterial and venous networks (Professional agreement) [15,16]. MRI with MR angiography can be performed as the first-intention exploration if compatible with the patient's clinical status and if it does not delay therapeutic management (Professional agreement) [16].

If subarachnoid hemorrhage is suspected and the imaging results do not provide evidence for the diagnosis, a lumbar puncture should always be performed, even if the headache has subsided (Grade B) [13,14,17]. CSF xanthochromia is present in 100% of patients with aneurismal subarachnoid hemorrhage when the CSF is drawn between 12 hr and 14 days after symptom onset and submitted to spectrophotometric analysis [5,13,18,19]. Some laboratories rely on a simple visual inspection of the supernatant that has lower sensitivity and specificity, creating a non-negligible risk of false-positive, and most importantly false-negative, results (Grade B) [20–22]. If the CT angiogram and the lumbar puncture do not provide evidence for subarachnoid hemorrhage related to a vascular malformation, cerebral angiography is not useful (Grade B) [17,23,24]. Subarachnoid hemorrhage without vascular malformation may correspond to a reversible cerebral vasoconstriction syndrome (RCVS). Other explorations may be discussed on a case-by-case basis, particularly if the pain persists.

In patients with suspected subarachnoid hemorrhage, if the CT angiography (MRA ou MR angiography) does not provide evidence for the diagnosis, a lumbar puncture should always be performed, even if the headache has subsided (Grade B) (Edlow et al., 2008; Steiner et al., 2013; Stewart et al., 2014).

A lumbar puncture using an atraumatic 25-gauge needle is recommended in order to reduce the risk of post-lumbar puncture headache and limit further induced costs (Grade B) (Davis et al., 2014; Lavi et al., 2006; Strupp et al., 2001; Tung et al., 2012).

There are many non-traumatic etiologies of sudden-onset headache. The most common causes are listed in Table 2. Other causes can also produce the same clinical presentation: CSF hypotension, angina pectoris or intra-ventricular tumor.

³ HAS Haute Autorité de Santé.

Table 2 – The most common etiologies of sudden-onset headache.

	Potentially associated clinical features	Complementary explorations
Subarachnoid hemorrhage	Meningeal syndrome Isolated third nerve palsy Loss of consciousness	Brain MRI + brain arterial angiogram sequences Brain CT ± CT brain angiogram ± lumbar puncture Arteriography
Other intracranial bleeding	Focal signs that can be discrete in certain localizations (cerebellum)	Brain MRI ± brain CT ± brain CT angiogram
Brain infarction	Focal signs that can be discrete in certain localizations (cerebellum)	Brain MRI + brain arterial angiogram sequences Brain CT ± CT brain angiogram
Reversible cerebral vasoconstriction syndrome (RCVS)	Repeated episodes occurring spontaneously or induced by exercise, Valsalva maneuver, before or with orgasm. Possible focal signs or epilepsy	Brain MRI + brain arterial angiogram sequences Brain CT ± CT brain angiogram ± lumbar puncture ± arteriography
Cerebral venous thrombus	Neurological deficit Epileptic seizure Focal signs, intracranial hypertension	Brain MRI + brain arterial angiogram sequences Brain CT ± CT venous angiogram
Cervical arterial dissection	Neck pain, Claude-Bernard-Horner sign, pulsating tinnitus	Brain CT + CT angiogram of the superior aortic trunks (SAT), Ultrasound-Doppler of the SAT, Brain MRI + MRI angiogram of the head and neck Lumbar puncture
Meningitis ± encephalitis	Fever, meningeal syndrome, cranial nerve involvement	Ocular fundus: papillary edema Brain MRI
Hypertensive encephalopathy and eclampsia, posterior reversible encephalopathy syndrome (PRES)	Headache followed by perturbed consciousness, focal deficit, seizures High blood pressure, 240/120 mmHg (lower if eclampsia)	
Pituitary necrosis	Vision disorders, oculomotor disorders	Brain MRI
Temporal arteritis	(sudden-onset rare) Age > 50 yr Impaired general condition	C-reactive protein: elevated Temporal artery biopsy

8.1.1.3. Diagnostic strategy.

For patients with subarachnoid hemorrhage, investigations should be performed to search for the cause, preferably with CT or MR angiography **Professional agreement**. Arteriography may be discussed on a case-by-case basis.

If the clinical presentation is a thunderclap headache, especially when there is no vascular malformation, the etiological search should focus on an RCVS [25] (**Professional agreement**). RCVS was considered to be rare, but is probably underdiagnosed. Its exact incidence is not known, but in tertiary centers, RCVS has been diagnosed in 45% of patients presenting with thunderclap headache [26]. RCVS is attributed to transient reversible abnormal regulation of cerebral arterial tone, which triggers multifocal diffuse vasoconstriction and vasodilatation, favored by vasoactive substances such as cannabis, cocaine, ecstasy, amphetamines, LSD, antidepressants (serotonin or serotonin and noradrenalin reuptake inhibitors), nasal decongestants, triptans and ergotamine. Onset is sudden, typically with thunderclap headache, often triggered by sexual activity or Valsalva maneuvers. RCVS can be complicated or associated with hemorrhage, ischemia or cerebral artery dissection [27]. Diagnosis requires demonstration of typical arterial anomalies on the CT or MRI angiogram. The first imaging exploration may be normal if performed early during the first 4–5 days after symptom onset. Anomalies reach a maximum 2–3 weeks after the first symptoms.

Thunderclap headache can also be the inaugural sign of cervical artery dissection, cerebral venous thrombosis, or

pituitary apoplexy, with a normal physical examination, CT, and CSF. Other explorations must be proposed when the CT angiogram of the supra-aortic trunks and the CSF analysis are normal; exploration of the cervical vessels and a brain MRI are needed [28]. The delay for obtaining these explorations will be discussed with a specialist (neurologist, neurosurgeon, neuroradiologist) on a case-by-case basis.

If no diagnosis is established after the lumbar puncture, the initial imaging will be completed by imaging of the supra-aortic trunks. If an RCVS is suspected, the imaging will be repeated a few days later (MRI with MRI angiogram or, if not available, CT angiogram).

8.1.2. Progressive unusual headache (onset or worsening within the last 7 days)

History taking and the physical examination (with ocular fundus) search for signs of intracranial hypertension (headache particularly intense at awakening in the morning, vomiting, visual blurring, papillary edema at the ocular fundus) that are suggestive of an expansive intracranial process or neck pain suggestive of dissection of the cervical arteries.

In case of unusual headache that came on or worsened within the last 7 days, brain imaging should be performed rapidly to search for an expansive intracranial process or a vascular cause (Professional agreement).

Ideally, brain MRI with T1, T2, FLAIR, and T1 injected sequences will be performed to search for contrast uptake and




Fig. 2 – Schematic representation of the diagnostic strategy. CT: computed tomography; CRP: C-reactive protein; MRI: magnetic resonance imaging; CSF: cerebral spinal fluid; RCVS: cerebral vasoconstriction syndrome; PRES: posterior reversible encephalopathy syndrome.

signs of venous thrombosis. A T2* sequence can also be ordered to identify potential bleeding and contribute to the search for venous thrombosis. The supra-aortic trunks will be explored to search for dissection, warranting a fat-saturation sequence but also MR angiography of the supra-aortic trunks (Professional agreement).

Unless there is an emergency, and if MRI is easily accessible, CT is not needed [29]. If MRI accessibility is limited, a brain CT without contrast injection should be performed (Professional agreement).

If intracranial hypertension is suspected and imaging does not provide the diagnosis, CSF analysis is needed (Professional agreement). Pressure should be measured in a specialized center. An ocular fundus is also needed.

If the imaging work-up fails to provide diagnostic evidence, lumbar puncture can be an option to search for meningitis or a CSF pressure disorder (Professional agreement). This puncture should not be made if spontaneous intracranial hypotension is suspected (orthostatic headache is part of the presentation). In this situation, brain MRI with injection should be preferred. Lumbar puncture will include a measure of opening pressure (hypotension < 6 cm H₂O, or hypertension > 25 cm H₂O) [5]. In the event of idiopathic intracranial hypertension, the ocular fundus visualizes papillary edema and the MRI may show an empty sella turcica, dilatation of the peri-optic subarachnoid spaces, narrowing of the lateral sinus, or a flatten posterior

portion of the sclera [30]. In the event of idiopathic intracranial hypotension, the cerebral MRI may show meningeal contrast uptake [31–33]. When intracranial hypotension is suspected, the MRI should be performed before the lumbar puncture (Professional agreement).

In patients aged over 50 years, blood tests will include cell counts, electrolytes, BUN, liver tests and C-reactive protein in all cases. An inflammatory syndrome is suggestive of giant-cell arteritis (Horton's disease) (Hunder *et al.*, 1990) (Professional agreement).

It is not recommended to assay D-dimers to rule out a diagnosis of cerebral venous thrombosis; approximately 20% of patients with thrombi causing isolated headache have a normal D-dimer level (Grade B) (Fig. 2).

8.2. Emergency care

The initial work-up should not only search for evidence of secondary headache, but should also identify the diagnostic framework in patients with primary headache. The description here will be limited to primary headache as care for secondary headache is determined by the etiology, associated with symptomatic treatment for pain relief. It is simply recalled that the treatment of RCVS is based on the administration of nimodipine, given empirically at the same dose as in subarachnoid hemorrhage, in combination with




Fig. 3 – Therapeutic management of acute migraine or status migrainosus in the emergency setting. In the event of nausea or vomiting, NSAID can be administered intramuscularly or intravenously, metoclopramide and hydroxyzine intravenously, and triptans by nasal spray or subcutaneously (sumatriptan). In general, intravenous administration should be preferred over oral administration; rehydration should be favored.

symptomatic treatment for headache (**Professional agreement**). All treatments with a vasoconstrictor effect should be discontinued. Furthermore, response to analgesic treatment is not indicative of a primary cause and should not be considered reassuring [17,34].

8.2.1. Migraine

For treatment of migraine attack, the patient should be advised to rest in a dark calm room (**Professional agreement**). Fluid replacement (oral or intravenous) is also important (**Professional agreement**). The medical treatment will be adapted to the patient's current regimen. Oral NSAID are the first-intention drugs (diclofenac 50–100 mg, ibuprofen 400 mg, ketoprofen 100–150 mg, naproxen 500 mg), or a combination of aspirin 900 mg + metoclopramide 10 g (**Grade A**) [35–38]. Ibuprofen and ketoprofen have marketing approval for migraine in France. It is recommended to use a dose of 400 mg for ibuprofen and 100–150 mg for ketoprofen. In patients with a contraindication for NSAID, one gram of acetaminophen combined with 10 mg metoclopramide can be a good alternative (**Grade A**) [39]. If the patient has already taken NSAIDs (or acetaminophen + metoclopramide) at a proper dose and without efficacy, triptans can be proposed (**Grade A**) [39]. For attacks that fail to respond to first-line treatment, metoclopramide in a slow intravenous infusion (10 mg/15 min) should always be given for its anti-emetic effect if the patient has nausea [37,40,41], but also for its anti-migraine effect [42,43] (**Grade A**). Since metoclopramide is a dopaminergic antagonist, an anti-cholinergic agent may be added to avoid extra-pyramidal effects, especially akathisia (**Grade C**). Slow administration avoids the risk of akathisia [44,45] (**Grade B**). In published studies, metoclopramide was combined with diphenhydramine (anti-histamine H1 agent

with anti-cholinergic properties), but this product is not available for intravenous administration in France. The closest product is hydroxyzine which we propose. It is however not common to use this product for this indication.

If the patient started by taking triptans, a NSAID may be prescribed (**Professional agreement**). If nausea and vomiting preclude oral intake, the NSAID (ketoprofen 100 mg, aspirin 1000 mg) may be administered by intravenous injection and metoclopramide by intravenous infusion [43,46–49] (**Grade B**). For triptans, the only product with a nasal spray formulation is sumatriptan. As a last resort, if the nasal spray is not tolerated, a subcutaneous injection can be used (**Grade A**) [50]. Intravenous infusion of amitriptyline (25 mg/2 h) can be proposed if migraine persists despite use of triptans and NSAIDs (**Professional agreement**). Dexamethasone (10 mg i.v.) can be used, despite moderate efficacy [51] (**Grade A**).

As a last resort, phenothiazine derivatives may be used. Prochlorperazine has a better level of proof, but is not available in France [52]. Otherwise, chlorpromazine (Largactil®) (slow i.v. infusion 0.1 mg/kg) can be proposed [53] (**Grade A**). The patient should be advised of the major sedative effect. Despite a good level of proof, the acute and chronic adverse effects of this treatment are non-negligible so that this therapeutic option should be strictly limited (**Professional agreement**).

It is recommended to not use magnesium intravenously (**Grade A**) [54]. It is also recommended to not use opiates (**Grade C**) [29,55,56].

8.2.2. Status migrainosus

There is no recognized treatment for this rare entity that is a diagnosis of exclusion. The patient should be referred to an emergency ward to ensure intravenous treatment. Amitriptyline can be used (i.v. 25 mg/2 hr) (**Professional agreement**).

Dexamethasone (i.v. 10 mg) can be used, despite moderate efficacy [51] (Grade A). Otherwise, chlorpromazine (Largactil[®]) (slow i.v. infusion 0.1 mg/kg) can be proposed [53] (Professional agreement). The patient should be advised of the major sedative effect. Despite a good level of proof, the acute and chronic adverse effects of this treatment are non-negligible so that this therapeutic option should be strictly limited (Professional agreement) (Fig. 3).

8.2.3. Tension-type headache

First-intension treatment relies on use of an analgesic drug (acetaminophen 1 g) or a NSAID (ibuprofen 400 mg or ketoprofen 100 mg) [57,58] (Grade A). Here again, metoclopramide can be added for its analgesic effect, with the usual precautions, if the attack fails to respond [59] (Grade A). As a last resort, amitriptyline can be proposed by intravenous infusion (25 mg) (Professional agreement).

8.2.4. Cluster headache

Two first-line treatments have marketing approval in France for acute attacks [7]:

- sumatriptan, 6 mg sc SC (Grade A);
- oxygen (12–15 L/min) using a high-concentration mask for 15–20 min (Professional agreement).

Disclosure of interest

X. Moisset reports non-financial support from Merck-Serono, Biogen, Sanofi-Pasteur-MSD, Genzyme and Roche; personal fees from Astellas and Institut UPSA de la douleur; outside the submitted work.

J. Mawet reports non-financial support from SOS Oxygene, non-financial support from AMGEN, personal fees and non-financial support from Air Liquide, outside the submitted work.

E. Guegan-Massardier reports non-financial support from Merck Serono, TEVA, Genzyme, BMS, Bayer Health Care SAS, LFB, Pfizer SAS, Biogen, outside the submitted work.

E. Bozzolo has nothing to disclose.

V. Gilard has nothing to disclose.

E. Tollard has nothing to disclose.

T. Feraud has nothing to disclose.

B. Noëlle reports non-financial support from Merck-Serono, Biogen, Novartis; personal fees from UCB; outside the submitted work.

C. Rondet has nothing to disclose.

A. Donnet reports support from Allergan, Almirall SAS, Amgen, Astellas, Elli Lilly & Compagny, Menarini, MSD, Pfizer Inc, Saint-Jude, Teva and Zambon, outside the submitted work.

Acknowledgements

We thank the members of the reading group: the French club against cluster headache (AFCAVF⁴), Alchaar Haiel (neurologist),

⁴ AFCAVF: Association française contre l'algie vasculaire de la face.

Alibeu Jean Pierre (anesthetist), Autret Alain (neurologist), Barège-Négre Michèle (neurologist), Brion Nathalie (therapist), De Diego Emmanuelle (general practitioner), De Gaalon Solène (neurologist), Derouiche Fayçal (neurologist), Domingo Valérie (neurologist), Dousset Virginie (neurologist), Ducros Anne (neurologist), Eschalière Bénédicte (general practitioner), Fabre Nelly (neurologist), Fontaine Gwladys (general practitioner), Fontaine Denys (neurosurgeon), Hinzelin Grégoire (neurologist), Kassnasrallah Saad (neurologist), Lajoie Jean-Louis (general practitioner), Lantéri-Minet Michel (neurologist), Lauer Valérie (neurologist), Leclerc Xavier (neuroradiologist), Le Jeunne Claire (internist), Levraut Jacques (intensivist), Moustafa Fares (intensivist), Piano Virginie (algologist), Roos Caroline (neurologist), Salvat Eric (algologist) and Serrie Alain (anesthetist).

REFERENCES

- [1] Rasmussen BK, Jensen R, Schroll M, Olesen J. Epidemiology of headache in a general population—a prevalence study. *J Clin Epidemiol* 1991;44:1147–57.
- [2] Burch RC, Loder S, Loder E, Smitherman TA. The prevalence and burden of migraine and severe headache in the United States: updated statistics from government health surveillance studies. *Headache* 2015;55:21–34. <http://dx.doi.org/10.1111/head.12482>.
- [3] Goldstein JN, Camargo CA, Pelletier AJ, Edlow JA. Headache in United States emergency departments: demographics, work-up and frequency of pathological diagnoses. *Cephalalgia* 2006;26:684–90. <http://dx.doi.org/10.1111/j.1468-2982.2006.01093.x>.
- [4] Pari E, Rinaldi F, Gipponi S, Venturelli E, Liberini P, Rao R, et al. Management of headache disorders in the Emergency Department setting. *Neurol Sci Off J Ital Neurol Soc Ital Soc Clin Neurophysiol* 2015. <http://dx.doi.org/10.1007/s10072-015-2148-7>.
- [5] Headache Classification Committee of the International Headache Society (IHS). The International Classification of Headache Disorders, 3rd edition (beta version). *Cephalalgia* 2013;33:629–808. <http://dx.doi.org/10.1177/0333102413485658>.
- [6] Lantéri-Minet M, Demarquay G, Alchaar H, Bonnin J, Cornet P, Douay X, et al. Management of chronic daily headache in migraine patients: medication overuse headache and chronic migraine. French guidelines (French Headache Society, French Private Neurologists Association, French Pain Society). *Rev Neurol (Paris)* 2014;170:162–76. <http://dx.doi.org/10.1016/j.neurol.2013.09.006>.
- [7] Donnet A, Demarquay G, Ducros A, Geraud G, Giraud P, Guegan-Massardier E, et al. French guidelines for diagnosis and treatment of cluster headache (French Headache Society). *Rev Neurol (Paris)* 2014;170:653–70. <http://dx.doi.org/10.1016/j.neurol.2014.03.016>.
- [8] Beltramone M, Donnet A. Status migrainosus and migraine aura status in a French tertiary-care center: an 11-year retrospective analysis. *Cephalalgia* 2014;34:633–7. <http://dx.doi.org/10.1177/0333102414520763>.
- [9] Linn FH, Rinkel GJ, Algra A, van Gijn J. Headache characteristics in subarachnoid haemorrhage and benign thunderclap headache. *J Neurol Neurosurg Psychiatry* 1998;65:791–3.
- [10] Bø SH, Davidsen EM, Gulbrandsen P, Dietrichs E. Acute headache: a prospective diagnostic work-up of patients admitted to a general hospital. *Eur J Neurol* 2008;15:1293–9. <http://dx.doi.org/10.1111/j.1468-1331.2008.02279.x>.

- [11] Bellolio MF, Hess EP, Gilani WI, VanDyck TJ, Ostby SA, Schwarz JA, et al. External validation of the Ottawa subarachnoid hemorrhage clinical decision rule in patients with acute headache. *Am J Emerg Med* 2015;33:244–9. <http://dx.doi.org/10.1016/j.ajem.2014.11.049>.
- [12] Perry JJ, Stiell IG, Sivilotti MLA, Bullard MJ, Hohl CM, Sutherland J, et al. Clinical decision rules to rule out subarachnoid hemorrhage for acute headache. *JAMA* 2013;310:1248–55. <http://dx.doi.org/10.1001/jama.2013.278018>.
- [13] Steiner T, Juvola S, Unterberg A, Jung C, Forsting M, Rinkel G, et al. European Stroke Organization guidelines for the management of intracranial aneurysms and subarachnoid haemorrhage. *Cerebrovasc Dis Basel Switz* 2013;35:93–112. <http://dx.doi.org/10.1159/000346087>.
- [14] Stewart H, Reuben A, McDonald J. LP or not LP, that is the question: gold standard or unnecessary procedure in subarachnoid haemorrhage? *Emerg Med J* 2014;31:720–3. <http://dx.doi.org/10.1136/emered-2013-202573>.
- [15] Connolly ES, Rabinstein AA, Carhuapoma JR, Derdeyn CP, Dion J, Higashida RT, et al. Guidelines for the management of aneurysmal subarachnoid hemorrhage: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke J Cereb Circ* 2012;43:1711–37. <http://dx.doi.org/10.1161/STR.0b013e3182587839>.
- [16] Edjlali M, Rodriguez-Régent C, Hodel J, Aboukais R, Trystram D, Pruvo J-P, et al. Subarachnoid hemorrhage in ten questions. *Diagn Interv Imaging* 2015;96:657–66. <http://dx.doi.org/10.1016/j.diii.2015.06.003>.
- [17] Edlow JA, Panagos PD, Godwin SA, Thomas TL, Decker WW, American College of Emergency Physicians. Clinical policy: critical issues in the evaluation and management of adult patients presenting to the emergency department with acute headache. *Ann Emerg Med* 2008;52:407–36. <http://dx.doi.org/10.1016/j.annemergmed.2008.07.001>.
- [18] Al-Shahi R, White PM, Davenport RJ, Lindsay KW. Subarachnoid haemorrhage. *BMJ* 2006;333:235–40. <http://dx.doi.org/10.1136/bmj.333.7561.235>.
- [19] Vivancos J, Gilo F, Frutos R, Maestre J, García-Pastor A, Quintana F, et al. Clinical management guidelines for subarachnoid haemorrhage. Diagnosis and treatment. *Neurol Barc Spain* 2014;29:353–70. <http://dx.doi.org/10.1016/j.nrl.2012.07.009>.
- [20] Chadwick C, van Heyningen C. Visual inspection does not reliably detect bilirubin in CSF. *BMJ* 2013;346:f647.
- [21] Cruickshank A, Auld P, Beetham R, Burrows G, Egner W, Holbrook I, et al. Revised national guidelines for analysis of cerebrospinal fluid for bilirubin in suspected subarachnoid haemorrhage. *Ann Clin Biochem* 2008;45:238–44. <http://dx.doi.org/10.1258/acb.2008.007257>.
- [22] Hann A, Chu K, Greenslade J, Williams J, Brown A. Benefit of cerebrospinal fluid spectrophotometry in the assessment of CT scan negative suspected subarachnoid haemorrhage: a diagnostic accuracy study. *J Clin Neurosci Off J Neurosurg Soc Australas* 2015;22:173–9. <http://dx.doi.org/10.1016/j.jocn.2014.07.025>.
- [23] Landtblom A-M, Fridriksson S, Boivie J, Hillman J, Johansson G, Johansson I. Sudden onset headache: a prospective study of features, incidence and causes. *Cephalalgia* 2002;22:354–60.
- [24] Savitz SI, Levitan EB, Wears R, Edlow JA. Pooled analysis of patients with thunderclap headache evaluated by CT and LP: is angiography necessary in patients with negative evaluations? *J Neurol Sci* 2009;276:123–5. <http://dx.doi.org/10.1016/j.jns.2008.09.019>.
- [25] Ducros A. Reversible cerebral vasoconstriction syndrome. *Lancet Neurol* 2012;11:906–17. [http://dx.doi.org/10.1016/S1474-4422\(12\)70135-7](http://dx.doi.org/10.1016/S1474-4422(12)70135-7).
- [26] Cheng Y-C, Kuo K-H, Lai T-H. A common cause of sudden and thunderclap headaches: reversible cerebral vasoconstriction syndrome. *J Headache Pain* 2014;15:13. <http://dx.doi.org/10.1186/1129-2377-15-13>.
- [27] Mawet J, Boukobza M, Franc J, Sarov M, Arnold M, Bousser M-G, et al. Reversible cerebral vasoconstriction syndrome and cervical artery dissection in 20 patients. *Neurology* 2013;81:821–4. <http://dx.doi.org/10.1212/WNL.0b013e3182a2cbe2>.
- [28] Ducros A, Bousser M-G. Thunderclap headache. *BMJ* 2013;346:e8557.
- [29] Loder E, Weizenbaum E, Frishberg B, Silberstein S, American Headache Society Choosing Wisely Task Force. Choosing wisely in headache medicine: the American Headache Society's list of five things physicians and patients should question. *Headache* 2013;53:1651–9. <http://dx.doi.org/10.1111/head.12233>.
- [30] Peng K-P, Fuh J-L, Wang S-J. High-pressure headaches: idiopathic intracranial hypertension and its mimics. *Nat Rev Neurol* 2012;8:700–10. <http://dx.doi.org/10.1038/nrneurol.2012.223>.
- [31] Urbach H. Intracranial hypotension: clinical presentation, imaging findings, and imaging-guided therapy. *Curr Opin Neurol* 2014;27:414–24. <http://dx.doi.org/10.1097/WCO.000000000000105>.
- [32] Crassard I, Soria C, Tzourio C, Woimant F, Drouet L, Ducros A, et al. A negative D-dimer assay does not rule out cerebral venous thrombosis: a series of seventy-three patients. *Stroke J Cereb Circ* 2005;36:1716–9. <http://dx.doi.org/10.1161/01.STR.0000173401.76085.98>.
- [33] Dentali F, Squizzato A, Marchesi C, Bonzini M, Ferro JM, Ageno W. D-dimer testing in the diagnosis of cerebral vein thrombosis: a systematic review and a meta-analysis of the literature. *J Thromb Haemost JTH* 2012;10:582–9. <http://dx.doi.org/10.1111/j.1538-7836.2012.04637.x>.
- [34] Pope JV, Edlow JA. Favorable response to analgesics does not predict a benign etiology of headache. *Headache* 2008;48:944–50. <http://dx.doi.org/10.1111/j.1526-4610.2008.01127.x>.
- [35] Kirthi V, Derry S, Moore RA. Aspirin with or without an antiemetic for acute migraine headaches in adults. *Cochrane Database Syst Rev* 2013;4:CD008041. <http://dx.doi.org/10.1002/14651858.CD008041.pub3>.
- [36] Law S, Derry S, Moore RA. Naproxen with or without an antiemetic for acute migraine headaches in adults. *Cochrane Database Syst Rev* 2013;10:CD009455. <http://dx.doi.org/10.1002/14651858.CD009455.pub2>.
- [37] Marmura MJ, Silberstein SD, Schwedt TJ. The acute treatment of migraine in adults: the american headache society evidence assessment of migraine pharmacotherapies. *Headache* 2015;55:3–20. <http://dx.doi.org/10.1111/head.12499>.
- [38] Rabbie R, Derry S, Moore RA. Ibuprofen with or without an antiemetic for acute migraine headaches in adults. *Cochrane Database Syst Rev* 2013;4:CD008039. <http://dx.doi.org/10.1002/14651858.CD008039.pub3>.
- [39] Derry S, Moore RA. Paracetamol (acetaminophen) with or without an antiemetic for acute migraine headaches in adults. *Cochrane Database Syst Rev* 2013;4:CD008040. <http://dx.doi.org/10.1002/14651858.CD008040.pub3>.
- [40] Kelley NE, Tepper DE. Rescue therapy for acute migraine, part 1: triptans, dihydroergotamine, and magnesium. *Headache* 2012;52:114–28. <http://dx.doi.org/10.1111/j.1526-4610.2011.02062.x>.
- [41] Lanteri-Minet M, Valade D, Géraud G, Lucas C, Donnet A, Société française d'étude des migraines et des céphalées. Guidelines for the diagnosis and management of migraine in adults and children. *Rev Neurol (Paris)* 2013;169:14–29. <http://dx.doi.org/10.1016/j.neurol.2012.07.022>.

- [42] Friedman BW, Corbo J, Lipton RB, Bijur PE, Esses D, Solorzano C, et al. A trial of metoclopramide vs sumatriptan for the emergency department treatment of migraines. *Neurology* 2005;64:463-8. <http://dx.doi.org/10.1212/01.WNL.0000150904.28131.DD>.
- [43] Friedman BW, Garber L, Yoon A, Solorzano C, Wollowitz A, Esses D, et al. Randomized trial of IV valproate vs metoclopramide vs ketorolac for acute migraine. *Neurology* 2014;82:976-83. <http://dx.doi.org/10.1212/WNL.000000000000223>.
- [44] Parlak I, Atilla R, Cicek M, Parlak M, Erdur B, Guryay M, et al. Rate of metoclopramide infusion affects the severity and incidence of akathisia. *Emerg Med J* 2005;22:621-4. <http://dx.doi.org/10.1136/emj.2004.014712>.
- [45] Tura P, Erdur B, Aydin B, Turkcuier I, Parlak I. Slow infusion metoclopramide does not affect the improvement rate of nausea while reducing akathisia and sedation incidence. *Emerg Med J* 2012;29:108-12. <http://dx.doi.org/10.1136/emj.2010.094367>.
- [46] Gungor F, Akyol KC, Kesapli M, Celik A, Karaca A, Bozdemir MN, et al. Intravenous dexketoprofen vs placebo for migraine attack in the emergency department: A randomized, placebo-controlled trial. *Cephalalgia* 2015. <http://dx.doi.org/10.1177/0333102415584604>.
- [47] Taggart E, Doran S, Kokotillo A, Campbell S, Villa-Roel C, Rowe BH. Ketorolac in the treatment of acute migraine: a systematic review. *Headache* 2013;53:277-87. <http://dx.doi.org/10.1111/head.12009>.
- [48] Turkcuier I, Serinken M, Eken C, Yilmaz A, Akdag Ö, Uyan E, et al. Intravenous paracetamol versus dexketoprofen in acute migraine attack in the emergency department: a randomised clinical trial. *Emerg Med J* 2014;31:182-5. <http://dx.doi.org/10.1136/emered-2013-203044>.
- [49] Weatherall MW, Telzerow AJ, Cittadini E, Kaube H, Goadsby PJ. Intravenous aspirin (lysine acetylsalicylate) in the inpatient management of headache. *Neurology* 2010;75:1098-103. <http://dx.doi.org/10.1212/WNL.0b013e3181f39a11>.
- [50] Derry CJ, Derry S, Moore RA. Sumatriptan (all routes of administration) for acute migraine attacks in adults - overview of Cochrane reviews. *Cochrane Database Syst Rev* 2014;5:CD009108. <http://dx.doi.org/10.1002/14651858.CD009108.pub2>.
- [51] Woldeamanuel Y, Rapoport A, Cowan R. The place of corticosteroids in migraine attack management: a 65-year systematic review with pooled analysis and critical appraisal. *Cephalalgia* 2015. <http://dx.doi.org/10.1177/0333102414566200>.
- [52] Orr SL, Aubé M, Becker WJ, Davenport WJ, Dilli E, Dodick D, et al. Canadian Headache Society systematic review and recommendations on the treatment of migraine pain in emergency settings. *Cephalalgia* 2014. <http://dx.doi.org/10.1177/0333102414535997>.
- [53] Kelly A-M, Walczynski T, Gunn B. The relative efficacy of phenothiazines for the treatment of acute migraine: a meta-analysis. *Headache* 2009;49:1324-32. <http://dx.doi.org/10.1111/j.1526-4610.2009.01465.x>.
- [54] Choi H, Parmar N. The use of intravenous magnesium sulphate for acute migraine: meta-analysis of randomized controlled trials. *Eur J Emerg Med Off J Eur Soc Emerg Med* 2014;21:2-9. <http://dx.doi.org/10.1097/MEJ.0b013e3283646e1b>.
- [55] Friedman BW, Grosberg BM. Diagnosis and management of the primary headache disorders in the emergency department setting. *Emerg Med Clin North Am* 2009;27. <http://dx.doi.org/10.1016/j.emc.2008.09.005>. 71-87, viii.
- [56] McCarthy LH, Cowan RP. Comparison of parenteral treatments of acute primary headache in a large academic emergency department cohort. *Cephalalgia* 2014. <http://dx.doi.org/10.1177/0333102414557703>.
- [57] Bendtsen L, Evers S, Linde M, Mitsikostas DD, Sandrini G, Schoenen J, et al. EFNS guideline on the treatment of tension-type headache - report of an EFNS task force. *Eur J Neurol* 2010;17:1318-25. <http://dx.doi.org/10.1111/j.1468-1331.2010.03070.x>.
- [58] Moore RA, Derry S, Wiffen PJ, Straube S, Bendtsen L. Evidence for efficacy of acute treatment of episodic tension-type headache: methodological critique of randomised trials for oral treatments. *Pain* 2014;155:2220-8. <http://dx.doi.org/10.1016/j.pain.2014.08.009>.
- [59] Weinman D, Nicastrò O, Akala O, Friedman BW. Parenteral treatment of episodic tension-type headache: a systematic review. *Headache* 2014;54:260-8. <http://dx.doi.org/10.1111/head.12287>.